

Quick Guidelines to Reviewing your Proof PDF below

1. Please mark your corrections in the proof PDF OR clearly describe corrections in a text file.

DO NOT: implement desired changes in the PDF or supply newly edited Word files, TEX files, or XML files.

2. When marking the PDF, clearly show or describe requested changes. We must be able to read every correction in order to add changes to our master copy of the proof (see examples below).

Example of correctly showing text deletion

Rhesus macaques are an ideal host species to study the behavioral bases of ~~dyadic-and/or community-wide~~ bacterial sharing. They are biologically, socially and cognitively analogous to human societies (*Cobb, 1976; Suomi, 2011*). In nature, they live in large (approximate range: 20–150 individuals), multi-male-multi-female social groups, in which individuals

Example of correctly showing text replacement

mice. Significance of *SIRT2* for TRIG in LCL was also replicated in GLGC at $P = 1.1 \times 10^{-102}$ (Table 5). The similar effects and significance of *SIRT2* in knockout mice studies, the Cebu cohort analysis, and GLGC PrediXcan results indicate the increased expression levels. Therefore, variation in the regulation of IG levels.

heather Reply X

that

1/23/2018 5:05 PM Post

nes associated with TRIG at the 11q23.3 locus, ng. This analysis showed that *BACE1* has the *MA1*, and *CEP164* contributing smaller effects.

st Asian populations

f our study have been previously associated ot conduct a robust gene-based association 2; *Rasmussen-Torvik et al., 2012*). In our 2 (*PAX6*) in Yoruba and additional genes at the chromosome 11 locus in Cebu, including genes with predictive SNPs located hundreds

2. Please check the entire article and **review all critical information carefully**. Examples ([this list is not comprehensive](#)):

- Title
- Names and email addresses of corresponding author(s)
- Author names & affiliations
- Figures
- Tables

3. Please answer all queries listed at the end of the article.

NOTES

We ask that you return proofs in 24 hours but **please do not rush**. Take the time that you need to review this thoroughly and submit all your corrections in a complete draft. **We only accept one set of corrections**.

Any substantive changes introduced at this stage such as the removal or addition of new text, revising Supplemental Information files, and changing table or figure values **requires Academic Editor approval**. This may cause delays in publication.

Please only
ANNOTATE
the proof.
Do not edit
the PDF.
If multiple
authors will
review this PDF,
please return
one file
containing all
corrections.

Submitted 11 April 2018
Accepted 29 August 2018
Published 16 October 2018

Corresponding author
Sidarta Ribeiro,
sidartaribeiro@neuro.ufrn.br

Academic editor
Stephen Macknik

Additional Information and
Declarations can be found on
page 9

DOI 10.7717/peerj.5678

© Copyright
2018 Pereira et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Non-visual exploration of novel objects increases the levels of plasticity factors in the rat primary visual cortex

Catia M. Pereira^{1,*}, Marco A.M. Freire^{2,*}, José R. Santos³, Joanielson S. Guimarães⁴, Gabriella Dias-Flores⁵, Sharlene Santos⁵, Antonio Pereira⁶ and Sidarta Ribeiro⁵

¹Instituto Internacional de Neurociências de Natal Edmond e Lily Safra, Macaíba, RN, Brasil

²Laboratório de Neurologia Experimental, Universidade Estadual do Rio Grande do Norte, Mossoró, RN, Brasil

³Departamento de Biociências, Universidade Federal de Sergipe, Itabaiana, SE, Brasil

⁴Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém, PA, Brasil

⁵Instituto do Cérebro, Universidade Federal do Rio Grande do Norte, Natal, RN, Brasil

⁶Faculdade de Engenharia Elétrica, Universidade Federal do Pará, Belém, PA, Brasil

*These authors contributed equally to this work.

ABSTRACT

Background. Historically, the primary sensory areas of the cerebral cortex have been exclusively associated with the processing of a single sensory modality. Yet, the presence of tactile responses in the primary visual (V1) cortex has challenged this view, leading to the notion that primary sensory areas engage in cross-modal processing, and that the associated circuitry is modifiable by such activity. To explore this notion, here we assessed whether the exploration of novel objects in the dark induces the activation of plasticity markers in the V1 cortex of rats.

Methods. Adult rats were allowed to freely explore for 20 min a completely dark box with four novel objects of different shapes and textures. Animals were euthanized either 1 ($n = 5$) or 3 h ($n = 5$) after exploration. A control group ($n = 5$) was placed for 20 min in the same environment, but without the objects. Frontal sections of the brains were submitted to immunohistochemistry to measure protein levels of *egr-1* and *c-fos*, and phosphorylated calcium-dependent kinase (pCaKMII), in V1 cortex.

Results. The amount of neurons labeled with monoclonal antibodies against *c-fos*, *egr-1* or pCaKMII increased significantly in V1 cortex after one hour of exploration in the dark. Three hours after exploration, the number of labeled neurons decreased to basal levels.

Conclusions. Our results suggest that non-visual exploration induces the activation of immediate-early genes in V1 cortex, which is suggestive of cross-modal processing in this area. Besides, the increase in the number of neurons labeled with pCaKMII may signal a condition promoting synaptic plasticity.

Subjects Neuroscience

Keywords Immediate-early gene, CaMKII, Phosphorylation, Cross-modal processing, Visual cortex

INTRODUCTION

At the beginning of the last century, Korbinian Brodmann proposed that the neocortex is divided into several cytoarchitecturally distinct areas, each one devoted to a distinct function (Brodmann, 2006; Guimaraes, Santos & Freire, 2016). This classical finding was subsequently reinforced by several studies using multiple experimental techniques, including tract-tracing studies showing that thalamocortical inputs from a given sensory modality preferentially target specific primary cortical areas (Hubel & Wiesel, 1959; Mountcastle, 1997; Pereira et al., 2000; Rocha et al., 2007; Santiago et al., 2007; Dooley et al., 2015). However, a growing number of recent studies have shown that neuronal responses in those areas are characterized by a large amount of sensory crosstalk (Lemus et al., 2010; Vincis & Fontanini, 2016). For instance, whisker stimulation in rats generates patterns of subthreshold cortical activation that spread far from the traditional limits of the primary somatosensory cortex (Frostig et al., 2008). This pattern of activation means that unimodal sensory stimulation can potentially modulate the activity of neurons located in sensory areas related to other sensory modalities, despite the existence of cytoarchitectural borders (Stehberg, Dang & Frostig, 2014).

It remains to be determined, however, whether such cross-modal interactions would impact beyond the transitory modulation of neuronal activity, to trigger longer-lasting plastic changes in primary sensory cortices. Monocular enucleation during adulthood elicits cross-modal reorganization of the visual cortex, a phenomenon supported by preexisting projections from the somatosensory cortex (Van Brussel, Gerits & Arckens, 2011). Thus, cross-modal plasticity may help individuals recover from sensory deprivation due to injury of the neural pathways associated with a specific modality (Rabinowitch & Bai, 2016).

In a previous study, we showed that the whisker-based exploration of novel objects in the darkness leads to a marked upregulation of the mRNA levels of calcium-dependent immediate-early genes (IEGs) in primary visual (V1) cortex of calcium-dependent immediate-early genes (IEGs) which are associated with memory consolidation (Ribeiro et al., 2007). Using multielectrode recordings and the same free-exploration paradigm, we also showed in a later study that 35% of the V1 neurons significantly change their firing rates during whisker-based object exploration, despite the lack of visual inputs (Vasconcelos et al., 2011). Most importantly, neuronal ensembles in V1 cortex were shown to carry non-visual information useful to discriminate among different objects (Vasconcelos et al., 2011).

As a follow up to these earlier studies, in the present work we sought to contribute with additional information concerning the modulation of visual cortex induced by tactile stimulation. We investigated the causes and consequences of cross-modal mRNA IEG induction in V1 cortex, by assessing the levels of phosphorylated CaMKII, a calcium-dependent kinase upstream of *c-fos* and *egr-1* transcripts with a well-documented role in plasticity (Kaczmarek & Chaudhuri, 1997). To examine the response downstream, we assessed the levels of *c-fos* and *egr-1* proteins (Bading, Ginty & Greenberg, 1993; Kaczmarek, 2000; Jones et al., 2001; Nunes et al., 2010).




Figure 1 **Experimental design.** Experimental animals were transferred to a box and allowed to freely explore four novel objects of different shapes and textures ('ball', 'urchin', 'brush' and 'food') in the dark during 20 min, being returned to their regular cages after the end of exploration. (B) Control animals were placed for 20 min in the same box, but without the objects. Drawing adapted from *Ribeiro et al. (2007)*.

Full-size DOI: [10.7717/peerj.5678/fig-1](https://doi.org/10.7717/peerj.5678/fig-1)

MATERIALS AND METHODS

Animals

Fifteen adult male Wistar rats (5 months-old, 320 ± 20 g) were used in this research. All experimental procedures were approved by the Institutional Local Ethics Committee for the Use of Animals (ID 05/2007), in accordance with the NIH Guidelines for the Care and Use of Laboratory Animals. All efforts were made to avoid suffering and to reduce the number of animals used. Prior to the experiment, all animals remained in conventional cages ($49 \times 34 \times 16$ cm) under standard conditions (lights on at 07:00, lights off at 19:00, 22 ± 2 °C) with free access to food and water.

Object exploration in the dark

All experimental procedures began at 22:00 under infrared light, to ensure that the animals were kept unstimulated by visible light for 3 h. A group of animals ($n = 10$) was transferred to a closed box ($55 \times 40 \times 20$ cm) and allowed to freely explore four novel objects of different shapes and textures (termed 'ball', 'urchin', 'brush' and 'food') for 20 min (Fig. 1A). The animals were then returned to their home cages and euthanized after one ($n = 5$) or three ($n = 5$) hours after exploration. A control group of animals ($n = 5$) was placed for 20 min in the same box in the dark, but without the objects (Fig. 1B), returned to their home cages and then euthanized after 1 h. Home cages were free of objects and the animals were well adapted to them, which precluded that prior environmental stimuli could induce IEG activation. To evaluate the degree of locomotion of the animals during the post-exploratory period, the total distance traveled and the mean velocity were video recorded and quantified using Any-maze software (version 4.3, Stoelting Co., Wood Dale, IL, USA).

Immunohistochemistry

All animals were deeply anaesthetized with sodium pentobarbital (90 mg/kg, i.p.) and decapitated. The brains were quickly removed and fast-frozen in an embedding medium (Tissue Tek, Sakura Finetek, Japan). Then, the brains were frontally sectioned at $20 \mu\text{m}$ in a

cryostat (Carl Zeiss Micron HM 550, Germany) and thaw-mounted over electrically charged glass slides following a serial distribution. Brain slices were then post-fixed for 15 min in 4% paraformaldehyde in 0.1 M phosphate buffer (PB). For immunohistochemistry, sections were washed during 20 min in phosphate buffer-Tween (PB-T) and incubated in a blocking buffer (BB) solution (0.5% fresh skim milk and 0.3% Triton X-100 in 0.1M PB) for 30 min. Sections were incubated overnight at 18 °C in primary antibody (1:200 in BB; phosphorylated polyclonal rabbit CaMKII, Millipore Corp., USA, catalog #AB3827; rabbit polyclonal egr-1 and c-fos, Santa Cruz Biotechnology, Inc., USA, catalog #sc-189 and #sc-52, respectively), washed in PB-T (2×, 10 min each), incubated with a biotinylated secondary goat anti-rabbit antibody (1:200 in BB, Vector Labs, USA) for 2 h, washed in PB-T (2×, 10 min each), and incubated in avidin-biotin-peroxidase solution (Vectastain Standard ABC kit, Vector Labs, USA) for 2 h. Next, slides were placed in a solution containing 0.03% 3,3' diaminobenzidine (DAB) and 0.001% hydrogen peroxide in 0.1 M PB (Freire *et al.*, 2005). Primary antibodies were replaced by normal serum in some test sections. The reaction was monitored every 30 min and was interrupted by washing sections in 0.1 M PB during 5 min. In order to avoid artifacts due to procedural inconsistencies, all sections of all groups were processed together, incubated simultaneously and spent an identical time in the same DAB solution. Alternate sections were stained with cresyl violet (Nissl staining) to reveal cyto-architectonic boundaries. At the end of all procedures, sections were dehydrated and cover-slipped with Entellan (Merck, Darmstadt, Germany). Digital images were acquired with a CX9000 camera (MBF Bioscience Inc., Williston, VT, USA) attached to a Nikon Eclipse 80i optical microscope (Tokyo, Japan—4×, 10× and 20×objectives) under the same light parameters, to avoid bias. The contrast, and/or brightness of pictures were adjusted using Photoshop CS5 software (Adobe Systems Inc., San José, CA, USA).

Data quantification

IEG-labeled cells (3 sections/animal, $n = 5$ animals per group for every marker) were counted with the *NeuroLucida* system (MBF Bioscience Inc., USA) using 10 grids of $100 \times 100 \mu\text{m}$ for each section across V1 cortex. Analysis of pCaMKII staining in V1 cortex was done by optical densitometry with the ImageJ software (<http://rsb.info.nih.gov/ij/>) using a 0.05 mm^2 square window ($n = 5$ animals per group, 3 sections per animal; 5 samples per section). The low reactivity in the white matter was used as background reference (signal was averaged over 5 different sites using the same square window). For each animal, the average optical density (OD) was named G , cortical white matter W and a contrast index C was calculated according to the equation: $C = (G - W) / (G + W)$ (Freire *et al.*, 2007). All photographs were taken with the same illumination and microscope (Nikon Eclipse 80i, Tokyo, Japan). Average values for all measurements were normalized to the levels observed in the control group.

Statistics

Significant differences across groups were assessed with the non-parametric Kruskal Wallis test, with Dunn *post-hoc* test ($\alpha = 0.05$), using the Graphpad Prism 5.0 software

(GraphPad Software Inc., La Jolla, CA, USA). The numerical values represented in boxplots were presented as median with indication of the 25th to 75th percentiles.

RESULTS

To ensure that the analyses involved only the region of interest associated with V1 cortex, its boundaries were defined by Nissl staining (Fig. 2A). In a qualitative analysis, immunohistochemistry with specific antibodies revealed a robust increase of *egr-1* and *c-fos* labeling in V1 cortex of animals killed 1 h after object exploration in darkness, with a subsequent decrease after 3 h (Fig. 2B). However, the pattern of activation at the 3 h time period was still higher than the basal levels found in unexposed controls (Fig. 2B). The levels of phosphorylated CaMKII in V1 cortex followed a similar time course (Fig. 2B).

Although we had not quantified their intensity of immunolabeling in *c-fos* and *egr-1*-reactive cells, the general pattern of reactivity allowed us to identify cell profiles varying from intensely to weakly reactive in all evaluated time points. There was a marked prevalence of intensely reactive cells 1 h after exploration, a pattern markedly different from that seen at 3 h, which was more balanced (Fig. 2B). The effectiveness of the immunolabeling method for all markers could be demonstrated by the absence of labeling when the primary antibody was replaced by normal serum (test sections). Since the brains were collected fresh and immediately fast-frozen, some blood normally remains in the tissue. This likely explains the differences in background labeling observed across animals in the present study.

A quantitative analysis confirmed that the peak of activity in V1 cortex occurred after 1 h of exploration in darkness for all markers (Fig. 3) (Kruskal–Wallis, Dunn *post-hoc* test, $*p < 0.05$). After 3 h of exploration the expression levels for all markers in V1 cortex were lower when compared to those seen at 1 h of exploration, being similar to control levels (Fig. 3). Locomotion parameters (total traveled distance and mean velocity) measured during the post-exploration period revealed a similar behavioral pattern in all groups (Fig. 4A), with the animals spending equivalent amounts of time moving in the home cage (Fig. 4B).

DISCUSSION

In the present work, we investigated whether the whisker-based exploration of novel objects in the dark is capable of increasing the protein levels of molecular markers of plasticity in the V1 cortex. In a previous study using the same experimental paradigm, we have demonstrated that it increases the mRNA levels of calcium-dependent IEGs in the V1 cortex (Ribeiro *et al.*, 2007). Here we provide evidence that such non-visual exploration also increases the levels of phosphorylated CaMKII, as well as *c-fos* and *egr-1* proteins. This suggests that the cross-modal response in the V1 cortex can lead to long-term plasticity.

In a recent study in rats, Nakajima and coworkers showed that visual deprivation induces a sustained increase of neural activity in the S1 cortex (Nakajima *et al.*, 2016). The up-regulation of neuronal responsiveness in the S1 cortex after visual deprivation can be explained by an strengthening of synaptic activity mediated by NMDA and/or AMPA receptors, which induces IEG expression (Kaczmarek & Chaudhuri, 1997). Our present




Figure 2 Transient activation of plasticity-related factors in V1 cortex after non-visual object exploration. (A) Drawing indicating the location and border V1/V2 and the general aspect of the brain, immediately prior to sectioning. (B) Nissl-stained frontal section showing V1 cortex, obtained from the region indicated with the black square in A. The dashed line indicates the border between V1 and V2. (C–K) Sections immunostained for *egr-1*, *c-fos* and pCaMKII. Controls were unexposed to the objects, and experimental animals were assessed 1 h and 3 h after new object exposure. Notice the increased labeling after 1 h of exposure for all markers. Black squares on low power pictures indicate location from where the enlarged pictures were obtained. Scale bars: 3 mm (A); 200 μ m (B); 100 μ m (C–K).

Full-size [DOI: 10.7717/peerj.5678/fig-2](https://doi.org/10.7717/peerj.5678/fig-2)

results showing IEG activation in V1 cortex after non-visual exploration in the dark are supported by a recent report of long-term spatiotemporal changes in *c-fos* and *arc* levels in this area after bilateral deafness in adult rats (Pernia et al., 2017). According to this study, the disruption of the auditory input induces thalamocortical reorganization with consequent modification in both auditory and visual cortical circuits, which leads to the overexpression of IEGs in V1 cortex and a further re-expression of these markers in the




Figure 3 Normalized levels plasticity-related factors in V1 cortex after non-visual object exploration. Group data for egr-1, c-fos and pCaMKII, normalized in each panel by the mean value across groups. There was a significant increase 1 h after object exploration in V1 cortex for all markers ($n = 5$ animals per group, median \pm quartiles of values for all brain sections assessed, Kruskal–Wallis, Dunn *post hoc* test, * $p < 0.05$ and ** $p < 0.01$ corrected for the number of comparisons).

Full-size [DOI: 10.7717/peerj.5678/fig-3](https://doi.org/10.7717/peerj.5678/fig-3)




Figure 4 Locomotion parameters during the post-novelty period. (A) Quantification of total distance traveled and mean velocity showed that animals of all groups spent similar amount of time moving within their home cages. Blue dot indicates start point, red dot indicates end point. (B) Representative examples of trajectories covered during the post-novelty period. Kruskal–Wallis, Dunn *post hoc* test, * $p > 0.05$ corrected for the number of comparisons.

Full-size [DOI: 10.7717/peerj.5678/fig-4](https://doi.org/10.7717/peerj.5678/fig-4)

auditory cortex after thalamocortical reordering (*Pernia et al., 2017*). A similar pattern of IEG activation was also verified after visual deprivation (*Takahata & Kaas, 2016*).

The antibodies used have been previously validated (*Blanco et al., 2015*). The cross-modal up-regulation of the protein levels of egr-1 and c-fos detected in the present study is congruent with the known induction kinetics of these IEG (*Herdegen et al., 1991; Herdegen & Leah, 1998; Lonergan et al., 2010*), likely in association with the differential expression of NMDA receptors (*Chaudhuri, 1997; Kaczmarek & Chaudhuri, 1997*).

The present results add to the notion that cross-modality can be induced by the transient deprivation of stimulus from a sensory modality (visual, in this case), i.e., without any damage to the sensory periphery (*Pascual-Leone & Hamilton, 2001; Merabet et al., 2008*). However, since we only evaluated factors involved in a rapid response (one or three hours after visual deprivation), more studies are required to comprehensively characterize this phenomenon downstream of the initial genomic response.

Human brain imaging studies have shown that the visual cortex is significantly activated when subjects close their eyes and imagine objects (*Kosslyn et al., 1995*). Such visual mental imagery occurs when “a visual short-term memory (STM) representation is present but the stimulus is not actually being viewed” (*Kosslyn & Thompson, 2003*). In rodents, mental imagery can influence subsequent encoding and recognition processes of landmarks (*Karimpur & Hamburger, 2018*). In the present study we cannot rule out that our experimental design triggers imagery in the rats. However, since they explored the objects in the dark without any prior visual presentation, remains to be determined whether visual images could be formed for objects never encountered before.

In a previous study with the same experimental design, electrophysiological and molecular changes in V1 cortex were interpreted as nonspecific effects of arousal (*Ribeiro et al., 2007*). Assuming this hypothesis as correct, V1 cortex responses during whisker-based exploration of novel objects in the dark would represent only a general alert signal, carrying no specific information about object identity. However, in a subsequent study we analyzed with more depth the electrophysiological responses of individual V1 and S1 neurons of rats submitted to the same paradigm of exploration in the dark (*Vasconcelos et al., 2011*). This study revealed no significant difference in V1 and S1 neuronal responses to the explored objects, and thus reinforced the proposal that V1 cortex is in fact recruited during the whisker-based exploration of objects in the dark. Further studies evaluating IEG expression in S1 cortex using a control group with anesthetized whiskers shall help draw a more complete picture of the cellular changes in primary cortical areas whisker-based exploration in the dark.

In face of the present results, one key question concerning activation of visual areas by tactile stimulation is: why does it occur? Primary sensory areas have been shown to prefer a given modality but to quickly engage in cross-modal processing depending on task demands, a property named metamodality (*Pascual-Leone & Hamilton, 2001*). The functional recruitment of the visual cortex for tactile processing in healthy adults after a short period of visual deprivation is believed to reflect the unmasking of pre-existing connections (*Merabet et al., 2008*).

The visual cortex can be activated by locomotion (*Dipoppa et al., 2018*). To assess whether this effect could explain the results we evaluated the locomotion of the animals during the post-novelty period. We found that all groups spent similar amounts of time moving within their home cages, which rules out non-specific activation induced by locomotion.

CONCLUSIONS

Our results show that even a short period of dark (3 h) is able to engage neurons in the V1 cortex in non-visual processing, leading to the up-regulation of molecular cascades that link sustained membrane depolarization with long-term gene regulatory changes. Further studies are required to a better understanding of the circuit dynamics associated with IEG activation following visual deprivation.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was supported by grants from Pew Latin-American Program in Biomedical Science, FINEP 01.06.1092.00, INCT-CNPq/MCT INCEMAQ 704134/2009, CNPq Universal 481506/2007-1, CAPES/SECYT 152/08. AASDAP –Brazil provided supplementary support. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Pew Latin-American Program in Biomedical Science: FINEP 01.06.1092.00.

INCT-CNPq/MCT INCEMAQ: 704134/2009.

CNPq Universal: 481506/2007-1.

CAPES/SECYT: 152/08.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Catia M. Pereira and Marco A.M. Freire conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- José R. Santos and Joailson S. Guimarães performed the experiments, approved the final draft.
- Gabriella Dias-Florencio performed the experiments, analyzed the data, prepared figures and/or tables, approved the final draft.
- Sharlene Santos performed the experiments, prepared figures and/or tables, approved the final draft.

All supplemental information will be made available for download exactly as they were supplied. This link to the SI will only work when the article is published.

- Antonio Pereira contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Sidarta Ribeiro conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

All experimental procedures were approved by the Institutional Local Ethics Committee for the Use of Animals (ID 05/2007), in accordance with the NIH Guidelines for the Care and Use of Laboratory Animals.

Data Availability

The following information was supplied regarding data availability:

The raw data are provided in a [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5678#supplemental-information>.

REFERENCES

- Bading H, Ginty DD, Greenberg ME. 1993.** Regulation of gene expression in hippocampal neurons by distinct calcium signaling pathways. *Science* **260**:181–186 DOI [10.1126/science.8097060](https://doi.org/10.1126/science.8097060).
- Blanco W, Pereira CM, Cota VR, Souza AC, Rennó-Costa C, Santos S, Dias G, Guerreiro AMG, Tort ABL, Neto AD, Ribeiro S. 2015.** Synaptic homeostasis and restructuring across the sleep-wake cycle. *PLOS Computational Biology* **11**(5):e1004241 DOI [10.1371/journal.pcbi.1004241](https://doi.org/10.1371/journal.pcbi.1004241).
- Brodmann K. 2006.** *Brodmann's localisation in the cerebral cortex the principles of comparative localisation in the cerebral cortex based on cytoarchitectonics. (Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des Zellenbaues. Leipzig: Johann Ambrosius Barth Verlag; 1909).* Garey L, trans. New York: Springer.
- Chaudhuri A. 1997.** Neural activity mapping with inducible transcription factors. *Neuroreport* **8**:iii–vii.
- Dipoppa M, Ranson A, Krumin M, Pachitariu M, Carandini M, Harris KD. 2018.** Vision and locomotion shape the interactions between neuron types in mouse visual cortex. *Neuron* **98**:602–615 DOI [10.1016/j.neuron.2018.03.037](https://doi.org/10.1016/j.neuron.2018.03.037).
- Dooley JC, Franca JG, Seelke AM, Cooke DF, Krubitzer LA. 2015.** Evolution of mammalian sensorimotor cortex: thalamic projections to parietal cortical areas in *Monodelphis domestica*. *Frontiers in Neuroanatomy* **8**:163 DOI [10.3389/fnana.2014.00163](https://doi.org/10.3389/fnana.2014.00163).

- Freire MAM, Franca JG, Picanço-Diniz CW, Pereira Jr A. 2005.** Neuropil reactivity, distribution and morphology of NADPH diaphorase type I neurons in the barrel cortex of the adult mouse. *Journal of Chemical Neuroanatomy* **30**:71–81 DOI [10.1016/j.jchemneu.2005.04.006](https://doi.org/10.1016/j.jchemneu.2005.04.006).
- Freire MAM, Oliveira RB, Picanço-Diniz CW, Pereira A. 2007.** Differential effects of methylmercury intoxication in the rat's barrel field as evidenced by NADPH diaphorase histochemistry. *Neurotoxicology* **28**:175–181 DOI [10.1016/j.neuro.2006.06.007](https://doi.org/10.1016/j.neuro.2006.06.007).
- Frostig RD, Xiong Y, Chen-Bee CH, Kvasnak E, Stehberg J. 2008.** Large-scale organization of rat sensorimotor cortex based on a motif of large activation spreads. *Journal of Neuroscience* **28**:13274–13284 DOI [10.1523/JNEUROSCI.4074-08.2008](https://doi.org/10.1523/JNEUROSCI.4074-08.2008).
- Guimaraes JS, Santos JR, Freire MAM. 2016.** Korbinian Brodmann: cortical maps and the cytoarchitecture of the brain. *Academic Journal of Suriname* **7**:617–620.
- Herdegen T, Kovary K, Leah J, Bravo R. 1991.** Specific temporal and spatial distribution of JUN, FOS, and KROX-24 proteins in spinal neurons following noxious transsynaptic stimulation. *Journal of Comparative Neurology* **313**:178–191 DOI [10.1002/cne.903130113](https://doi.org/10.1002/cne.903130113).
- Herdegen T, Leah JD. 1998.** Inducible and constitutive transcription factors in the mammalian nervous system: control of gene expression by Jun, Fos and Krox, and CREB/ATF proteins. *Brain Research. Brain Research Reviews* **28**:370–490 DOI [10.1016/S0165-0173\(98\)00018-6](https://doi.org/10.1016/S0165-0173(98)00018-6).
- Hubel DH, Wiesel TN. 1959.** Receptive fields of single neurones in the cat's striate cortex. *Journal de Physiologie* **148**:574–591 DOI [10.1113/jphysiol.1959.sp006308](https://doi.org/10.1113/jphysiol.1959.sp006308).
- Jones MW, Errington ML, French PJ, Fine A, Bliss TV, Garel S, Charnay P, Bozon B, Laroche S, Davis S. 2001.** A requirement for the immediate early gene Zif268 in the expression of late LTP and long-term memories. *Nature Neuroscience* **4**:289–296 DOI [10.1038/85138](https://doi.org/10.1038/85138).
- Kaczmarek L. 2000.** Gene expression in learning processes. *Acta Neurobiologiae Experimentalis* **60**:419–424.
- Kaczmarek L, Chaudhuri A. 1997.** Sensory regulation of immediate-early gene expression in mammalian visual cortex: implications for functional mapping and neural plasticity. *Brain Research Reviews* **23**:237–256 DOI [10.1016/S0165-0173\(97\)00005-2](https://doi.org/10.1016/S0165-0173(97)00005-2).
- Karimpur H, Hamburger K. 2018.** A rat in the sewer: how mental imagery interacts with object recognition. *PLOS ONE* **13**:e0194227 DOI [10.1371/journal.pone.0194227](https://doi.org/10.1371/journal.pone.0194227).
- Kosslyn SM, Thompson WL. 2003.** When is early visual cortex activated during visual mental imagery? *Psychological Bulletin* **129**:723–746 DOI [10.1037/0033-2909.129.5.723](https://doi.org/10.1037/0033-2909.129.5.723).
- Kosslyn SM, Thompson WL, Kim IJ, Alpert NM. 1995.** Topographical representations of mental images in primary visual cortex. *Nature* **378**:496–498 DOI [10.1038/378496a0](https://doi.org/10.1038/378496a0).
- Lemus L, Hernandez A, Luna R, Zainos A, Romo R. 2010.** Do sensory cortices process more than one sensory modality during perceptual judgments? *Neuron* **67**:335–348 DOI [10.1016/j.neuron.2010.06.015](https://doi.org/10.1016/j.neuron.2010.06.015).

- Lonergan ME, Gafford GM, Jarome TJ, Helmstetter FJ. 2010.** Time-dependent expression of Arc and zif268 after acquisition of fear conditioning. *Neural Plast* 2010:139891.
- Merabet LB, Hamilton R, Schlaug G, Swisher JD, Kiriakopoulos ET, Pitskel NB, Kauffman T, Pascual-Leone A. 2008.** Rapid and reversible recruitment of early visual cortex for touch. *PLOS ONE* 3:e3046 DOI 10.1371/journal.pone.0003046.
- Mountcastle VB. 1997.** The columnar organization of the neocortex. *Brain* 120(Pt 4):701–722 DOI 10.1093/brain/120.4.701.
- Nakajima W, Jitsuki S, Sano A, Takahashi T. 2016.** Sustained enhancement of lateral inhibitory circuit maintains cross modal cortical reorganization. *PLOS ONE* 11:e0149068 DOI 10.1371/journal.pone.0149068.
- Nunes ACL, Duarte RB, Sousa TB, Santos JR, Freire MAM, Costa MS. 2010.** Expression of the immediate-early gene egr-1 and Substance P in the spinal cord following locomotor training in adult rats. *Brain Research* 1345:125–136 DOI 10.1016/j.brainres.2010.05.041.
- Pascual-Leone A, Hamilton R. 2001.** The metamodal organization of the brain. *Progress in Brain Research* 134:427–445 DOI 10.1016/S0079-6123(01)34028-1.
- Pereira Jr A, Freire MAM, Bahia CP, Franca JG, Picanço-Diniz CW. 2000.** The barrel field of the adult mouse SmI cortex as revealed by NADPH-diaphorase histochemistry. *Neuroreport* 11:1889–1892 DOI 10.1097/00001756-200006260-00017.
- Pernia M, Estevez S, Poveda C, Plaza I, Carro J, Juiz JM, Merchan MA. 2017.** c-Fos and Arc/Arg3.1 expression in auditory and visual cortices after hearing loss: evidence of sensory crossmodal reorganization in adult rats. *Journal of Comparative Neurology* 525:2677–2689 DOI 10.1002/cne.24233.
- Rabinowitch I, Bai J. 2016.** The foundations of cross-modal plasticity. *Communicative & Integrative Biology* 9:e1158378 DOI 10.1080/19420889.2016.1158378.
- Ribeiro S, Shi X, Engelhard M, Zhou Y, Zhang H, Gervasoni D, Lin SC, Wada K, Lemos NA, Nicolelis MA. 2007.** Novel experience induces persistent sleep-dependent plasticity in the cortex but not in the hippocampus. *Frontiers in Neuroscience* 1:43–55 DOI 10.3389/neuro.01.1.1.003.2007.
- Rocha EG, Santiago LF, Freire MAM, Gomes-Leal W, Lent R, Houzel JC, Franca JG, Pereira Jr A, Picanço-Diniz CW. 2007.** Callosal axon arbors in the limb representations of the somatosensory cortex (SI) in the agouti (*Dasyprocta primnolopha*). *Journal of Comparative Neurology* 500:255–266 DOI 10.1002/cne.21167.
- Santiago LF, Rocha EG, Freire MAM, Lent R, Houzel JC, Picanço-Diniz CW, Pereira Jr A, Franca JG. 2007.** The organizational variability of the rodent somatosensory cortex. *Reviews in the Neurosciences* 18:283–294.
- Stehberg J, Dang PT, Frostig RD. 2014.** Unimodal primary sensory cortices are directly connected by long-range horizontal projections in the rat sensory cortex. *Frontiers in Neuroanatomy* 8:93.
- Takahata T, Kaas JH. 2016.** c-FOS expression in the visual system of tree shrews after monocular inactivation. *Journal of Comparative Neurology* 525:151–165.

Van Brussel L, Gerits A, Arckens L. 2011. Evidence for cross-modal plasticity in adult mouse visual cortex following monocular enucleation. *Cereb Cortex* **21**:2133–2146 DOI [10.1093/cercor/bhq286](https://doi.org/10.1093/cercor/bhq286).

Vasconcelos N, Pantoja J, Belchior H, Caixeta FV, Faber J, Freire MAM, Cota VR, Anibal de Macedo E, Laplagne DA, Gomes HM, Ribeiro S. 2011. Cross-modal responses in the primary visual cortex encode complex objects and correlate with tactile discrimination. *Proceedings of the National Academy of Sciences of the United States of America* **108**:15408–15413 DOI [10.1073/pnas.1102780108](https://doi.org/10.1073/pnas.1102780108).

Vincis R, Fontanini A. 2016. Associative learning changes cross-modal representations in the gustatory cortex. *Elife* **5**.

Q3

Author Queries

Journal: PEERJ
Article id: 5678
Author: Pereira et al.
Title: Non-visual exploration of novel objects increases the levels of plasticity factors in the rat primary visual cortex

Q1 (Page 1)

Please check all author names, initials, and affiliations to confirm they are accurate.

Q2 (Page 10)

References Dooley et al. (2015), Lonergan et al. (2010), Stehberg et al. (2014) appear with one page number instead of a page range. If these are not single page references, please provide the page ranges. If the journal only uses article numbers for citations instead of page ranges, please provide that article number. If any of these are abstracts, please confirm which are single page abstracts and we will insert “[Abstract]” in the reference(s).

Q3 (Page 13)

Reference Vincis & Fontanini (2016) is incomplete. Please provide the missing information: page range.