


Can polyphenols improve the gut health status in pre-clinical study with diet-induced obesity?

A protocol for systematic review and/or meta-analysis

Lêda Karla Monteiro Dias, BSc^a, Gidyenne Christine Bandeira Silva de Medeiros, PhD^b, Ana Karolinne Nascimento Silva, UG^b, Ana Heloneida de Araujo Morais, PhD^{a,b,c}, Juliana Kelly da Silva-Maia, PhD^{a,b,*} 

Abstract

Introduction: Obesity is characterized as a low-grade inflammation that impairs physiological functions, including intestinal functioning and gut microbiota balance. Dietary polyphenols can be a strategy for obesity management, collaborating to preserve or recover gut health through antioxidant and anti-inflammatory actions, as well as modulators of the microbiota. This study describes a systematic review protocol to elucidate effects of polyphenols on intestinal health of pre-clinical models with diet-induced obesity. AIM: Our aim is to evaluate evidence about polyphenols' effects in the gut microbiota composition and diversity, parameters of the physical and molecular status of the gut barrier in obese models, additionally, understand the possible involved mechanisms.

Methodology: A protocol was developed and published on PROSPERO (Registration No: CRD42021262445). Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols is used to outline the protocol. The articles will be selected according to the PICOS strategy (population, interventions, control, outcome, and study design) in the following databases: PubMed, Science Direct, Scopus, Web of Science, and EMBASE. Experimental studies performed on rats and mice with a control group that describes treatment with polyphenols (from food matrix or crude extracts or isolated compounds) at any frequency, time, and dose will be included. Two reviewers will, independently, select the papers, extract data, and evaluate the data quality. The Systematic Review Center for Laboratory Animal Experimentation (SYRCLE) tool will be used to assess the risk of bias.

Expected results: Results will be showed through of native synthesis and, if possible, a metanalysis will be conducted. The review produced with this protocol can show the scientific evidence level about polyphenols' effects in intestinal health in obesity status.

Abbreviations: CFU = colony-forming unit, MeSH = Medical Subject Headings, OTUs = Operational taxonomic units, PICOS = population, interventions, control, outcomes, and study design, PRISMA = Preferred Reporting Items for Systematic Reviews and Meta-Analyses, PRISMA-P = Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols, PROSPERO = International Prospective Register of Systematic Reviews, SFCA = Short-chain fatty acids, SYRCLE = Systematic Review Center for Laboratory Animal Experimentation.

Keywords: dysbiosis, gut health, microbiota, obesity, polyphenols

The authors have no conflicts of interest to disclose.

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

^a Nutrition Postgraduate Program, Health Sciences Center, Federal University of Rio Grande do Norte, Natal, Brazil, ^b Department of Nutrition, Federal University of Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil, ^c Biochemistry and Molecular Biology Postgraduate Program, Center for Biosciences, Federal University of Rio Grande do Norte, Natal, Brazil.

* Correspondence: Juliana Kelly da Silva-Maia, Nutrition, Health Sciences Center, Federal University of Rio Grande do Norte, Natal, Brazil (e-mail: juliana.maia@ufrn.br).

Copyright © 2021 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Dias LK, Medeiros GC, Silva AK, Morais AH, Silva-Maia JK. Can polyphenols improve the gut health status in pre-clinical study with diet-induced obesity? A protocol for systematic review and/or meta-analysis. *Medicine* 2021;00(00).<https://doi.org/10.1097/MD.00000000000028162>.

Received: 17 November 2021 / Accepted: 18 November 2021

<http://dx.doi.org/10.1097/MD.00000000000028162>

1. Introduction

Bioactive compounds, such as polyphenols, are found in different food sources and can modulate inflammation and oxidative stress by acting in metabolic vias from obesity and other noncommunicable diseases.^[1]

Polyphenols are secondary metabolites from vegetables with key roles as a defense against microorganisms, insects, and environmental stress.^[2] Furthermore, these compounds have therapeutic actions such as antimicrobial, antiproliferative, antioxidant, and anti-inflammatory, which can act in local and systemic sites.^[3,4] From those functional properties, polyphenols have been suggested as adjuvants in obesity treatment.^[5-7]

Nowadays, obesity is a major and growing public health problem worldwide. According to the World Health Organization, 39% of adults were overweight and 13% were obese.^[8] This is a multifactorial and complex disease with an excessive accumulation of adipose tissue. There is evidence of correlation among gut microbiota disbalance, physiopathology process, and metabolic disturbers present on obesity.^[9,10]

Intestinal dysbiosis can be associated with low-grade inflammation from obesity that induces significant modifications to the gut microbiome. Eutrophic individuals show more diverse gut microbiota than those with fat tissue accumulation, which can contribute to metabolic disturbs through an axis of communication with adipose tissue.^[11,12] The dominative gut microbial phyla are Firmicutes, Bacteroidetes, Actinobacteria, Proteobacteria, and changes in the diversity and abundance of microorganisms impact health. Changes in the gut microbiota composition in obese individuals have been found, with a higher ratio of *Firmicutes* to *Bacteroidetes* as well as a reduced diversity of microbiota in the gut compared with eutrophic individuals.^[13] The imbalance between pathogenic and nonpathogenic microorganisms in the gut impairs the epithelial barrier and intestinal functions, that is, cause lower nutrient absorption, higher permeability to bacteria cytokines, and induce inflammatory response.^[14]

Sequentially, gut microbiota unbalance impairs the production of short-chain fatty acids (SCFA), which are yielded by metabolization of fermentable and non-digested compounds as saccharides, including glycosylated phenolic compounds. Examples of bacteria species with this role are *Lactobacillus*, *Lachnospiraceae*, and *Ruminococcaceae*.^[15] SCFA are key to maintaining intestinal health because are a fuel source for intestine epithelial cells, preserving the gut barrier integrity, besides reducing the intraluminal pH that avoids the pathogenic bacteria growing and stimulating the mucus production by caliciform cells. Moreover, impaired intestinal barrier increases the risk of bacteria translocation and other harmful substances leading to the production of inflammatory cytokines and contributing to keeping chronic inflammations as in obesity.^[16]

Unbalanced diet composition can contribute to obesity perpetuation and intestinal health disturbs. On the other hand, a healthier diet can be a strategy to provide bioactive compounds, like polyphenols, protecting against diseases. Indeed, diet interventions have been broadly studied as determinant factors to intestinal health and microbiota modulation. Highlighting that intestinal health is referred, besides to microbiota balance, to mucosal barrier integrity and function, that is, preserved processes as digestion, absorption, and host defense.^[4] In this sense, polyphenols are possible therapeutic approaches to prevent and treat obesity and the associated dysbiosis, due to their capacity to modulate the gut microbiota and intestinal permeability, as well as control inflammatory and oxidative processing.^[17] Additionally, the microbiota composition can influence polyphenols' effects because the produced metabolites through these compounds depend, in part, on gut microbiota metabolism.^[13]

Preclinical experiments with animal models are important to elucidate the effects and involved mechanisms in different situations, which can be tricky in clinical experiments once required more invasive procedures as biopsies. Data from experimental studies bring much information that are essentials to clinical research advances.^[18] In this regard, a systematic review gathering data from preclinical studies that assessed the bioactive compounds' effects and mechanisms in obesity-associated dysbiosis and other parameters about health intestinal is necessary to point the scientific evidence level and direct the future clinical experiments and applications.^[16,17,19]

Therefore, the review that will be produced with this protocol aims to show the scientific evidence about polyphenols' effects in intestinal health in obesity status, and possible action mechanisms as agents with antioxidant, anti-inflammatory, and of balance microbiota properties.

2. Methods

2.1. Protocol and registration

The protocol was prepared according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols (PRISMA-P).^[20] This protocol was registered in the International Prospective Register of Systematic Reviews (PROSPERO) on August 27, 2021 with registration number CRD42021262445, and available at https://www.crd.york.ac.uk/prospéro/display_record.php?RecordID=262445. Ethical approval is not necessary for this research, because it is a review study with the use of secondary data.

The systematic review will follow the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).^[21] The flowchart was created to demonstrate the selection of study articles PRISMA (Fig. 1). The Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) checklist^[22] was applied to improve the quality of the systematic review data.

2.2. Review question

Can polyphenols improve the gut health status in pre-clinical studies with diet-induced obesity?

2.3. Eligibility criteria

Original articles must attend the eligibility criteria according to the PICOS (population, interventions, control, outcomes, and study design) (Table 1).

2.4. Inclusion criteria

This review will include experimental studies with rats or mice of both sexes and ages, evaluating the effect of polyphenols in the treatment of diet-induced obesity, regarding the action in the intestinal health. The models must be treated with isolated polyphenols or crude extracts with identified polyphenols or food matrix with identified polyphenols. All types of routes and doses of administration will be included.

2.5. Exclusion criteria

Documents that are not scientific articles, case reports, reviews, studies with other animal models. Studies without polyphenols identification or studies that combine treatment with polyphenols and antibiotics, probiotics, or prebiotics. Studies that do not time of the experiment, frequency, and doses administered, and studies without a control group. Studies that do not investigated effects in intestinal health will be excluded.

2.6. Search strategy

Searches will be conducted in the following databases: PubMed; Science Direct; Scopus; Web of Science; BVS and EMBASE by 2 independent researchers (L.K.M.D and A.K.N.S.). The full search strategy will be based on the search components: polyphenols OR flavonoids OR phenols OR anthocyanins; obesity OR adiposity; rats OR mice; microbiota OR microbiome OR disbiosis. The final search strategy will be tailored using Medical Subject Headings (MeSH) terms, entry terms, and synonyms linked with Boolean operators.




Figure 1. Article selection flowchart. Adapted from Preferred Reporting Items for Systematic Reviews.

2.7. Selection process

The list of references of all studies included in the systematic review will be screened. Before final analysis, the searches will be updated. No restrictions will be applied to language and year of publication, or geographic limits.

All articles by the search strategy must be exported to the Rayyan QCRI application (The Systematic Reviews Web).^[2,3] Duplicated studies will be removed, following, then, the initial selection by considering their titles, abstract, and keywords.

Discrepancies between the reviewers will be identified and resolved through consensus, and a third reviewer (J.K.SM) will be involved when necessary to decide whether to include the study. Studies excluded will be recorded, and their reason for exclusion will be reported in the review. All researchers will then review the full text of all studies considered eligible for inclusion. The management of references will be applied to the software Mendeley.^[2,4] In the first step of data synthesis, we will present the results of the study selection process using the PRISMA statement flowchart.^[2,5]

Table 1
Elements of the research question according to the PICOS strategy.

Description	Abbreviation	Elements of the question
Population	P	Rats (<i>Rattus norvegicus</i>) and mice (<i>Mus musculus</i>) of both sexes and ages with diet-induced obesity.
Interventions	I	Treatment with isolated polyphenols or extracts with identified polyphenols.
Control/comparison	C	Rats (<i>Rattus norvegicus</i>) and mice (<i>Mus musculus</i>) with diet-induced obesity without treatment.
Outcomes	O	Markers associated with gut health status and microbiota intestinal.
Study design	S	In vivo experimental studies

PICOS = problem, intervention, control, outcomes and study design.

Antioxidant and anti-inflammatory properties, as well as microbiota modulators, are properties from polyphenols, which increases the interest in these compounds as a strategy in obesity treatment.^[38–40] Moreover, the use of the conventional drugs to treat obesity is limited by no specificity and side effects of nowadays available medicines.^[41]

Polyphenols can be metabolized by gut bacteria stimulating the growth of beneficial species as *Lactobacillus* and *Bifidobacteria*, and reducing the pathogenic ones as coliforms.^[42] These compounds act also in control of the inflammatory pathways, reducing the proinflammatory cytokines as LPS, TLR, TNF, IL-1, and IL-1R.^[43] A recent study showed that Wistar rats fed with a high-fat diet and treated with grape marc had significant improvements since weight loss, control of glycemia and lipid, until better microbiota profile, with high ratio of *Lactobacillus* spp. and *Bacteroidetes* spp.^[44] Other work with blueberry has found similar results, showing gut microbiota modulation, improvement of barrier integrity (increased villousities and mucin and β -defensin-2 expression), attenuating obesity-associated inflammation (lower proinflammatory cytokines expression), and better response to insulin in rats with diet-induced obesity.^[45]

Therefore, it is suggested that polyphenols can be applied to the treatment of obesity-associated inflammatory processes and maintenance of intestinal health. However, literature data about polyphenols' effects on obesity and intestinal health are fragmented challenging to affirm their efficacy and action mechanisms. This way, it is necessary to realize a systematic review focusing on the polyphenols action on intestinal health concomitantly with obesity treatment. The present protocol will address a systematic review that can cover the aforementioned gaps, consequently, direct future works and application of polyphenols in obesity treatment.

Acknowledgments

This work was supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Finance Code 001 – CAPES),

Author contributions

Conceptualization: Lêda Karla Monteiro Dias, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Data curation: Lêda Karla Monteiro Dias, Gidyenne Christine Bandeira Silva de Medeiros, Ana Karolinne Nascimento Silva, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Formal analysis: Lêda Karla Monteiro Dias, Gidyenne Christine Bandeira Silva de Medeiros, Ana Karolinne Nascimento Silva, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Investigation: Lêda Karla Monteiro Dias, Gidyenne Christine Bandeira Silva de Medeiros, Ana Karolinne Nascimento Silva, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Methodology: Lêda Karla Monteiro Dias, Gidyenne Christine Bandeira Silva de Medeiros, Ana Karolinne Nascimento Silva, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Project administration: Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Resources: Juliana Kelly Silva-Maia.

Supervision: Gidyenne Christine Bandeira Silva de Medeiros, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Validation: Lêda Karla Monteiro Dias, Gidyenne Christine Bandeira Silva de Medeiros, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Writing – original draft: Lêda Karla Monteiro Dias, Gidyenne Christine Bandeira Silva de Medeiros, Ana Karolinne Nascimento Silva, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

Writing – review & editing: Lêda Karla Monteiro Dias, Gidyenne Christine Bandeira Silva de Medeiros, Ana Karolinne Nascimento Silva, Ana Heloneida de Araujo Morais, Juliana Kelly Silva-Maia.

References

- Oliveira CBC, de , Brito LA, et al. Obesity: inflammation and bioactive compounds Carla. *J Heal Biol Sci* 2020;8:1.
- Lyu X, Lee J, Chen WN. Potential natural food preservatives and their sustainable production in yeast: terpenoids and polyphenols. Published online 2019;doi:10.1021/acs.jafc.8b07141.
- Cardona F, Andrés-Lacueva C, Tulipani S, Tinahones FJ, Queipo-Ortuño MI. Benefits of polyphenols on gut microbiota and implications in human health. *J Nutr Biochem* 2013;24:1415–22.
- Bernardi S, Bo C, Del , Marino M, et al. Polyphenols and intestinal permeability: rationale and future perspectives. *J Agric Food Chem* 2020; doi:10.1021/acs.jafc.9b02283.
- Zhao Y, Chen B, Shen J, et al. The beneficial effects of quercetin, curcumin, and resveratrol in obesity. *Oxid Med Cell Longev* 2017;2017: 1459497.
- Novotny JA, Chen TY, Terekhov AI, et al. The effect of obesity and repeated exposure on pharmacokinetic response to grape polyphenols in humans. *Mol Nutr Food Res* 2017;61:1–6.
- Zhang N-N, Guo W-H, Hu H, et al. Effect of a polyphenol-rich canarium album extract on the composition of the gut microbiota of mice fed a high-fat diet. *Molecules* 2018;23:9.
- World Health Organization (WHO). Obesity and overweight. World Health Organization (WHO). 2021. Available at: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. Accessed in November 2021.
- Fitria O. Transplante Fecal E Sua Possível Aplicação No Tratamento Da Obesidade: Uma Revisão De Literatura the. *J Chem Inf Model* 2013;53: 1689–99.
- Muscogiuri G, Cantone E, Cassarano S, et al. Gut microbiota: a new path to treat obesity. *Int J Obes Suppl* 2019;9:10–9.
- Sun L, Ma L, Ma Y, Zhang F, Zhao C, Nie Y. Insights into the role of gut microbiota in obesity: pathogenesis, mechanisms, and therapeutic perspectives. *Protein Cell* 2018;9:397–403.
- Aron-wisnewsky J, Prifti E, Belda E, et al. Major microbiota dysbiosis in severe obesity: fate after bariatric surgery Published online 2019;70–82. doi:10.1136/gutjnl-2018-316103.
- Liu J, He Z, Ma N, Chen ZY. Beneficial effects of dietary polyphenols on high-fat diet-induced obesity linking with modulation of gut microbiota. *J Agric Food Chem* 2020;68:33–47.
- Alves BKR, Santos LC, Dos SOUSA PVDL, Santos GM, Dos Barros NVDA. Prevalência de Sinais e Sintomas Sugestivos de Disbiose Intestinal em Acadêmicos de Uma Instituição de Ensino Superior. *Rev Bras Obesidade Nutr e Emagrecimento* 2020;14:588–97.
- Alves-santos AM, Sandra C, Sugizaki A, Lima GC. Prebiotic effect of dietary polyphenols: a systematic review. *J Funct Foods* 2020;74 (September):104169.
- Verediano TA, Stampini Duarte Martino H, Dias Paes MC, Tako E. Effects of anthocyanin on intestinal health: a systematic review. *Nutrients* 2021;13:4.
- Peron G, Hidalgo-Liberona N, González-Domínguez R, et al. Exploring the molecular pathways behind the effects of nutrients and dietary polyphenols on gut microbiota and intestinal permeability: a perspective on the potential of metabolomics and future clinical applications. *J Agric Food Chem* 2020;68:1780–9.
- Bernardi L, Malinoski B, Cain JP, Antonovicz S, Amaral LA do. Experimental models for studies in nutrition: hypertension and diabetes mellitus. *Visão Acadêmica* 2017;17:69–87.
- Guirro M, Costa A, Gual-Grau A, et al. Effects from diet-induced gut microbiota dysbiosis and obesity can be ameliorated by fecal microbiota transplantation: a multiomics approach. *PLoS One* 2019;14:1–19.
- Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1177–85.

- [21] Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* 2021;372.
- [22] Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (prisma-p) 2015: Elaboration and explanation. *BMJ* 2015;349(January):1–25.
- [23] Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016;5:1–10.
- [24] Zaugg H, West RE, Tateishi I, Randall DL. Mendeley: creating communities of scholarly inquiry through research collaboration. *TechTrends* 2011;55:32–6.
- [25] Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *PLoS Med* 2021;18:1–15.
- [26] Sandoval-Ramírez BA, Catalán Ú, Pedret A, et al. Exploring the effects of phenolic compounds to reduce intestinal damage and improve the intestinal barrier integrity: a systematic review of in vivo animal studies. *Clin Nutr* 2021;40:1719–32.
- [27] Nash V, Ranadheera CS, Georgousopoulou EN, et al. The effects of grape and red wine polyphenols on gut microbiota: a systematic review. *Food Res Int* 2018;113:277–87.
- [28] Koch W. Dietary polyphenols—important non-nutrients in the prevention of chronic noncommunicable diseases. A systematic review. *Nutrition* 2019;11:5.
- [29] Gomes JVP, Rigolon TCB, Souza MS, da S, et al. Antiobesity effects of anthocyanins on mitochondrial biogenesis, inflammation, and oxidative stress: a systematic review. *Nutrition* 2019;66:192–202.
- [30] Ballard RC, Galvão FT, Cazarin BBC, Maróstica RM. Effects of polyphenol-rich fruit extracts on diet-induced obesity in rodents: systematic review and meta-analysis. *Curr Pharm Des* 2019;25:3484–97.
- [31] DeWeerd S. Microbiome: a complicated relationship status. *Nature* 2014;508(7496 SUPPL):8–10.
- [32] Li H, Lelliott C, Håkansson P, et al. Intestinal, adipose, and liver inflammation in diet-induced obese mice. *Metabolism* 2008;57:1704–10.
- [33] Castañer O, Godaya J, Park Y-M, et al. Response to: comment on “The Gut Microbiome Profile in Obesity: A Systematic Review”. *Int J Endocrinol* 2018;2018:1–9.
- [34] Gomes AC, Hoffmann C, Mota JF. The human gut microbiota: metabolism and perspective in obesity. *Gut Microbes* 2018;9:308–25.
- [35] Boulangé CL, Neves AL, Chilloux J, Nicholson JK, Dumas ME. Impact of the gut microbiota on inflammation, obesity, and metabolic disease. *Genome Med* 2016;8:1–12.
- [36] Covasa M. Gut microbiota modulates intestinal signalling in obesity. *Appetite* 2016;96:643.
- [37] Tavares S, Araújo C, Bressan J. Intestinal microbiota; relevance to obesity and modulation by prebiotics and probiotics. *Nutrición Hospitalaria* 2013;28:1039–48.
- [38] Fazel S, Luigi G, Daglia M, Mohammad S. Role of quercetin as an alternative for obesity treatment: You are what you eat!. *FOOD Chem* 2015;179:305–10.
- [39] Carpené C, Gomez-zorita S, Deleruyelle S, Carpené MA. Novel strategies for preventing diabetes and obesity complications with natural polyphenols. *Curr Medicinal Chem* 2015;150–64.
- [40] Bruno RS, Neilson AP, Lambert JD, Moustaid-Moussa N. Journal of Nutritional Biochemistry Special Issue: Polyphenols, obesity, and cardiometabolic health. *J Nutr Biochem* 2021;89:2020–2. doi:10.1016/j.jnutbio.2020.108565.
- [41] Bray GA, Heisel WE, Afshin A, et al. The science of obesity management: an endocrine society scientific statement. *Endocr Rev* 2018;39:79–132.
- [42] Tomás-Barberán FA, Andrés-Lacueva C. Polyphenols and health: current state and progress. *J Agric Food Chem* 2012;60:8773–5.
- [43] Corrêa TA, Rogero MM. Polyphenols regulating microRNAs and inflammation biomarkers in obesity. *Nutrition* 2019;59:150–7.
- [44] Gerardi G, Cavia-Saiz M, Rivero-Pérez MD, González-SanJosé ML, Muñoz P. Wine pomace product modulates oxidative stress and microbiota in obesity high-fat diet-fed rats. *J Funct Foods* 2020;68(March):1–10.
- [45] Lee S, Keirse KL, Kirkland R, Grunewald ZI, Fischer JG, de La Serre CB. Blueberry supplementation influences the gut microbiota, inflammation, and insulin resistance in high-fat-diet-fed rats. *J Nutr* 2018;148:209–19.